

SOLUTION OF FISHERS EQUATION USING HOMOTOPY PERTURBATION ABOODH TRANSFORM METHOD

OLUDAPO OMOTOLA OLUBANWO¹, JULIUS TEMIATYO ADEPOJU^{1*}, ABIODUN
SUFIAJ AJANI¹, SUNDAY SENAYON IDOWU¹

ABSTRACT. In this study, a nonlinear one dimensional reaction Diffusion Equation called Fisher's equation is solved using Aboodh transform coupled with Homotopy Perturbation Method (AHPM). This Method is applied to the general one dimensional Fishers equation. This technique is demonstrated with three unique examples with different initial conditions. The simplicity of the Aboodh transform contributes to its rapid convergence rate from approximate to exact solutions, rendering it both easy to use and reliable.

1. INTRODUCTION

Fisher's equation is a typical illustration of a PDE, which serves as one of the foundational model for understanding population dynamics and dispersion in a variety of biological and ecological contexts [26, 6, 18]. When examining how diffusion and local growth rates affect a population's long-term geographic distribution, this equation is crucial [19, 8].

$$\frac{\partial \phi}{\partial t} = \rho \nabla^2 \phi + k\phi(1 - \phi) \quad (1.1)$$

The population density is represented by ϕ in this equation, time is represented by t the diffusion coefficient by ρ , and the growth rate parameter by k . It exemplifies fundamental ideas in mathematical ecology and biology by showing how populations react to the interaction of geographic dispersion and local growth [9, 12].

Over time, numerous methods have been employed to solve and simplify partial differential equations, particularly those that are nonlinear. HAM [10], STDM [14], VIM [4, 5], HPM [5, 7, 17], LTDM [27], MHPM [22] HPMLT [15], AHPM [21], MAHPM [24] and LHPM [23] are a few examples. A powerful mathematical technique known as the Aboodh Homotopy Perturbation Method (AHPM) may be used to a wide range of complex differential equations in science and engineering [28, 16, 21, 1]. This methodology combines the homotopy perturbation

Date: Received: Mar 20, 2024; Accepted: Apr 6, 2024.

* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 46L55; Secondary 44B20.

Key words and phrases. He's Polynomial, Reaction Diffusion Equation, Partial Differential Equation, Homotopy Perturbation Method.

method , a perturbation-based approach to approximating solutions, with the Aboodh transform, which effectively linearizes differential equations. By combining these two approaches, AHPM offers a flexible and organized framework for managing nonlinear problems, giving specialists and researchers a helpful tool to handle challenging issues. This method simplifies the analysis of complex population dynamics and reveals how factors

2. MATERIALS AND METHODS

In this section, we consider a general nonlinear Klein-Gordon equation. We use Homotopy perturbation method, to decompose the nonlinear term, so that the solution can be obtained with iteration procedure.

2.1. Aboodh Transform. We employ a transform known as the Aboodh transform, developed to investigate functions in the set A , specifically for functions of exponential order. [1]

$$A = \{\phi(t) : \exists M, k_1, k_2 > 0 \mid |\phi(t)| < Me^{-ut}\} \quad (2.1)$$

Where $M < \infty \forall k_1, k_2 \leq \infty$. The Aboodh transform is defined as an integral equation of the form [1].

$$\mathcal{A}[\phi(t)] = R(u) = \frac{1}{u} \int_0^\infty \phi(t) e^{-ut} dt \quad t \geq 0 \quad k_1 \leq u \leq k_2 \quad (2.2)$$

TABLE 1. The Aboodh transform of some partial differential functions [1]

$\mathcal{A}[\phi(y, t)]$	$\psi(y, u)$
$\mathcal{A} \left[\frac{\partial \phi(x, t)}{\partial y} \right]$	$R'(y, u)$
$\mathcal{A} \left[\frac{\partial^2 \phi(y, t)}{\partial y^2} \right]$	$R''(y, u)$
$\mathcal{A} \left[\frac{\partial^n \phi(y, t)}{\partial y^n} \right]$	$R^{(n)}(y, u)$
$\mathcal{A} \left[\frac{\partial \phi(y, t)}{\partial t} \right]$	$vR(y, u) - \frac{\phi(y, 0)}{u}$
$\mathcal{A} \left[\frac{\partial^2 \phi(y, t)}{\partial t^2} \right]$	$= v^2 R(y, u) - \frac{1}{u} \frac{\partial \phi(y, 0)}{\partial t} - \phi(y, 0)$

TABLE 2. Aboodh transform of Some Functions [1]

$\phi(t)$	$K_f \{\phi(t)\} = \psi(u)$
1	$\frac{1}{u^2}$
t	$\frac{1}{u^3}$
t^n	$\frac{n!}{u^{2n+1}}$
e^{at}	$\frac{1}{u^2 - au}$
$\sin(at)$	$\frac{a}{u(u^2 + a^2)}$
$\cos(at)$	$\frac{1}{u^2 + a^2}$

3. ABOODH HOMOTOPY PERTURBATION METHOD (AHPM)

The general partial differential equation is of the form [25]:

$$L\phi(y, t) + R\phi(y, t) + N\phi(y, t) = f(y, t) \quad (3.1)$$

L	Linear Operator
R	Coefficient of the Laplace operator
N	Coefficient of Nonlinear Term
$f(y, t)$	Source term

Taking the Aboodh transform of both sides of equation (3.1),

$$\mathcal{A}\{L\phi(y, t)\} = \mathcal{A}\{f(y, t)\} - \mathcal{A}\{R\phi(y, t) + N\phi(y, t)\}, \quad (3.2)$$

$$u\mathcal{A}\{\phi(y, t)\} - \frac{\phi(y, 0)}{u} = \mathcal{A}\{f(y, t)\} - \mathcal{A}\{R\phi(y, t) + N\phi(y, t)\}, \quad (3.3)$$

$$u\mathcal{A}\{\phi(y, t)\} = \frac{\phi(y, 0)}{u} + \mathcal{A}\{f(y, t)\} - \mathcal{A}\{R\phi(y, t) + N\phi(y, t)\} \quad (3.4)$$

Multiply equation (3.4) by $\frac{1}{u}$

$$\mathcal{A}\{\phi(y, t)\} = \frac{1}{u} \left[\frac{\phi(y, 0)}{u} + \mathcal{A}\{f(y, t)\} \right] - \frac{1}{u} \mathcal{A}\{R\phi(y, t) + N\phi(y, t)\} \quad (3.5)$$

Taking the Aboodh inverse transform of (3.5), we obtain

$$\mathcal{A}^{-1}\{\mathcal{A}\{\phi(y, t)\}\} = \mathcal{A}^{-1}\left\{\frac{1}{u} \left[\frac{\phi(y, 0)}{u} + \mathcal{A}\{f(y, t)\} \right]\right\} - \mathcal{A}^{-1}\left\{\frac{1}{u} \mathcal{A}\{R\phi(y, t) + N\phi(y, t)\}\right\} \quad (3.6)$$

Let $G(y, t) = \mathcal{A}^{-1}\left\{\frac{1}{u} \left[\frac{\phi(y, 0)}{u} + \mathcal{A}\{f(y, t)\} \right]\right\}$. Then, we have

$$\phi(y, t) = G(y, t) - \mathcal{A}^{-1}\left\{\frac{1}{u} \mathcal{A}\{R\phi(y, t) + N\phi(y, t)\}\right\} \quad (3.7)$$

Since the approximation is independent of t , so all its derivatives with respect to t are zero. Then, we obtain

$$\phi_{r+1}(y, t) = G(y, t) - \mathcal{A}^{-1}\left\{\frac{1}{u} \mathcal{A}\{R\phi_m(y, t) + N\phi_m(y, t)\}\right\} \quad (3.8)$$

Applying Homotopy Perturbation Method to equation (3.8). Assume equation (3.8) can also be expressed as

$$\begin{aligned} \phi_m(y, t) &= \phi_0(y, t) + p\phi_1(y, t) + p^2\phi_2(y, t) + p^3\phi_3(y, t) + \dots \\ \phi(y, t) &= \sum_{r=0}^{\infty} p^r \phi_r(y, t) \end{aligned} \quad (3.9)$$

Taking into consideration the nonlinear term, it can be simplified as

$$N\phi(y, t) = \sum_{r=0}^{\infty} p^r H_r(\phi) \quad (3.10)$$

Where H_m is the polynomial ϕ that denotes the nonlinear terms, and $H_m(\phi)$. The initial terms are given by

$$\begin{aligned} H_0(\phi) &= \phi_0\phi_0 \\ H_1(\phi) &= \phi_0\phi_1 + \phi_1\phi_0 \\ H_2(\phi) &= \phi_0\phi_2 + \phi_1\phi_1 + \phi_2\phi_0 \\ H_3(\phi) &= \phi_0\phi_3 + \phi_1\phi_2 + \phi_2\phi_1 + \phi_3\phi_0 \end{aligned} \quad (3.11)$$

Then, by substituting (3.9) and (3.10) into (3.8), we have

$$\sum_{m=0}^{\infty} p^m \phi_{r+1}(y, t) = G(y, t) + p \left(\mathcal{A}^{-1} \left\{ \frac{1}{u} \mathcal{A} \left(\sum_{m=0}^{\infty} R p^m \phi_{ry} + \sum_{m=0}^{\infty} H_m(\phi) \right) \right\} \right) \quad (3.12)$$

$$\begin{aligned} \phi_0 + p^1 \phi_1 + p^2 \phi_2 + \dots &= G(y, t) - p \mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \left\{ \left(\sum_{m=0}^{\infty} R \phi_{0y} + \sum_{r=0}^{\infty} H_0(\phi) \right) \right. \right. \\ &\quad \left. \left. + p \left(\sum_{m=0}^{\infty} R \phi_{1y} + \sum_{m=0}^{\infty} H_1(\phi) \right) + p^2 \left(\sum_{m=0}^{\infty} R \phi_{2y} + \sum_{m=0}^{\infty} H_2(\phi) \right) + \dots \right\} \right] \end{aligned} \quad (3.13)$$

The desired approximation is obtained by equating the corresponding exponents of p in equation (3.13) such that $\phi_0(y, t)$ is defined as $\phi_0(y, t) = \alpha + t\beta$, where α and β denotes the initial conditions.

$$p^0 : \phi_0(y, t) = G(y, t) = \mathcal{A}^{-1} \left\{ \frac{1}{u} \left[\frac{\phi(y, 0)}{u} + \mathcal{A} \{ f(y, t) \} \right] \right\} \quad (3.14)$$

$$p^1 : \phi_1(y, t) = \mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \left\{ \left(\sum_{r=0}^{\infty} R \phi_{0y} + \sum_{r=0}^{\infty} H_0(\phi) \right) \right\} \right] \quad (3.15)$$

$$p^2 : \phi_2(y, t) = \mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \left\{ \left(\sum_{m=0}^{\infty} R \phi_{1y} + \sum_{m=0}^{\infty} H_1(\phi) \right) \right\} \right] \quad (3.16)$$

$$p^3 : \phi_3(y, t) = \mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \left\{ \left(\sum_{m=0}^{\infty} R \phi_{2y} + \sum_{m=0}^{\infty} H_2(\phi) \right) \right\} \right] \quad (3.17)$$

4. MAIN RESULT

Here, nonlinear one dimensional Fishers Equation will be solved using AHPM

Example 4.1. Considering the one dimensional Fishers type equation of the form [2]

$$\phi_t - \phi_{yy} - 1(1 - \phi)\phi = 0 \quad (4.1)$$

With

$$\phi(y, 0) = \mu \quad (4.2)$$

Solution

We can rewrite equation (4.1) as

$$\frac{\partial \phi}{\partial t} - \frac{\partial^2 \phi}{\partial y^2} - 1(1 - \phi)\phi = 0 \quad (4.3)$$

$$\frac{\partial \phi}{\partial t} = \frac{\partial^2 \phi}{\partial y^2} + \phi - \phi^2 \quad (4.4)$$

Taking the Aboodh transform of equation (4.4),

$$\mathcal{A}\left\{\frac{\partial \phi}{\partial t}\right\} = \mathcal{A}\left\{\frac{\partial^2 \phi}{\partial y^2} + \phi - \phi^2\right\} \quad (4.5)$$

$$u\mathcal{A}\{\phi(y, t)\} - \frac{\phi(y, 0)}{u} = \mathcal{A}\left\{\frac{\partial^2 \phi}{\partial y^2} + \phi - \phi^2\right\} \quad (4.6)$$

$$u\mathcal{A}\{\phi(y, t)\} = \frac{\phi(y, 0)}{u} + \mathcal{A}\left\{\frac{\partial^2 \phi}{\partial y^2} + \phi - \phi^2\right\} \quad (4.7)$$

Multiply equation (4.7) by $\frac{1}{u^2}$

$$\mathcal{A}\{\phi(y, t)\} = \frac{1}{u} \left[\frac{\phi(y, 0)}{u} \right] + \frac{1}{u} \mathcal{A}\left\{\frac{\partial^2 \phi}{\partial y^2} + \phi - \phi^2\right\} \quad (4.8)$$

Taking the inverse Aboodh transform of equation (4.8), we obtain

$$\mathcal{A}^{-1}\{\mathcal{A}\{\phi(y, t)\}\} = \mathcal{A}^{-1}\left\{\frac{1}{u} \left[\frac{\phi(y, 0)}{u} \right]\right\} + \mathcal{A}^{-1}\left[\frac{1}{u} \mathcal{A}\left\{\frac{\partial^2 \phi}{\partial y^2} + \phi - \phi^2\right\}\right] \quad (4.9)$$

Since $\phi(y, 0) = \mu$. Then, we have

$$\phi(y, t) = \mathcal{A}^{-1}\left\{\frac{\mu}{u}\right\} - \mathcal{A}^{-1}\left[\frac{1}{u} \mathcal{A}\left\{\frac{\partial^2 \phi}{\partial y^2} + \phi - \phi^2\right\}\right] \quad (4.10)$$

Since the approximation is independent of t , so all its derivatives with respect to t are zero. Then, we obtain

$$\phi_{r+1}(y, t) = \mu t - \mathcal{A}^{-1}\left[\frac{1}{u} \mathcal{A}\left\{\frac{\partial^2 \phi}{\partial y^2} + \phi - \phi^2\right\}\right] \quad (4.11)$$

Applying Homotopy Perturbation Method to equation (4.11). Assume equation can be written as

$$\phi(y, t) = \phi_0(y, t) + p\phi_1(y, t) + p^2\phi_2(y, t) + p^3\phi_3(y, t) + \dots \quad (4.12)$$

$$\phi(y, t) = \sum_{m=0}^{\infty} p^m \phi_m(y, t) \quad (4.13)$$

Taking into consideration the nonlinear term, it can be simplified as

$$N\phi(y, t) = \sum_{m=0}^{\infty} p^m H_m(\phi) \quad (4.14)$$

Where H_m is the polynomial ϕ that denotes the nonlinear terms, and $H_m(\phi)$. The initial terms are given by

$$\begin{aligned} H_0(\phi) &= \phi_0\phi_0 \\ H_1(\phi) &= \phi_0\phi_1 + \phi_1\phi_0 \\ H_2(\phi) &= \phi_0\phi_2 + \phi_1\phi_1 + \phi_2\phi_0 \\ H_3(\phi) &= \phi_0\phi_3 + \phi_1\phi_2 + \phi_2\phi_1 + \phi_3\phi_0 \end{aligned}$$

Then, by substituting (4.13) and (4.14) into (4.11) we have

$$\sum_{m=0}^{\infty} p^m \phi_{m+1}(y, t) = \mu t + p \left(\mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \left\{ \sum_{m=0}^{\infty} \frac{\partial^2 \phi_m}{\partial y^2} + \sum_{m=0}^{\infty} \phi_m - \sum_{m=0}^{\infty} H_m(\phi) \right\} \right] \right) \quad (4.15)$$

$$\begin{aligned} \phi_0 + p^1 \phi_1 + p^2 \phi_2 + \dots &= \mu - p^0 \mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \left\{ \left(\sum_{m=0}^{\infty} \frac{\partial^2 \phi_0}{\partial y^2} + \sum_{m=0}^{\infty} \phi_0 - \sum_{m=0}^{\infty} H_0^2(\phi) \right) \right. \right. \\ &\quad + p^1 \left(\sum_{m=0}^{\infty} \frac{\partial^2 \phi_1}{\partial y^2} + \sum_{m=0}^{\infty} \phi_1 - \sum_{m=0}^{\infty} H_1^2(\phi) \right) \\ &\quad \left. \left. + p^2 \left(\sum_{m=0}^{\infty} \frac{\partial^2 \phi_2}{\partial y^2} + \sum_{m=0}^{\infty} \phi_2 - \sum_{m=0}^{\infty} H_2^2(\phi) \right) + \dots \right\} \right] \quad (4.16) \end{aligned}$$

The desired approximation is obtained by equating the corresponding exponents of p in equation (4.16) such that $\phi_0(y, t)$ is defined as $\phi_0(y, t) = \alpha + t\beta$, where α and β denotes the initial conditions.

$$p^0 : \phi_0(y, t) = \mu \quad (4.17)$$

$$p^1 : \phi_1(y, t) = \mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \left\{ \left(\frac{\partial^2 \phi_0}{\partial y^2} + \phi_0 - H_0^2(\phi) \right) \right\} \right] \quad (4.18)$$

$$= \mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \{ (\mu - \mu^2) \} \right]$$

$$= \mathcal{A}^{-1} \left[\frac{1}{u} \cdot \left[\frac{\mu}{u^2} - \frac{\mu^2}{u^2} \right] \right]$$

$$= \mathcal{A}^{-1} \left[\frac{\mu}{u^3} - \frac{\mu^2}{u^3} \right]$$

$$= (\mu - \mu^2) t$$

$$p^2 : \phi_2(y, t) = \mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \left\{ \left(\frac{\partial^2 \phi_1}{\partial y^2} + \phi_1 - H_1^2(\phi) \right) \right\} \right] \quad (4.19)$$

$$= \mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \{ (\mu t - \mu^2 t - \mu(\mu t - \mu^2 t) - \mu(\mu t - \mu^2 t)) \} \right]$$

$$= \mathcal{A}^{-1} \left[\frac{1}{u} \mathcal{A} \{ (\mu t - \mu^2 t - \mu^2 t - \mu^2 t) \} \right]$$

$$= (\mu - 3\mu^2 + 2\mu^3) \frac{t^2}{2!} \quad (4.20)$$

The approximate solution to equation (4.1) is given as

$$\sum_{r=0}^{\infty} \phi_r(y, t) = \mu + (\mu - \mu^2) t + (\mu - 3\mu^2 + 2\mu^3) \frac{t^2}{2!} + \dots \quad (4.21)$$

The exact solution of equation (4.21) is written as

$$\phi(y, t) = \frac{\mu e^t}{1 - \mu + \mu e^t} \quad (4.22)$$

Equation (4.34) is the same as the solution obtained in [2]

Example 4.2. Let solve a nonlinear Fisher's equation of the form [25]:

$$\frac{\partial \phi(y, t)}{\partial t} = \frac{\partial^2 \phi(y, t)}{\partial y^2} + 6(1 - \phi(y, t))\phi(y, t) \quad (4.23)$$

With

$$\phi(y, 0) = e^y \quad (4.24)$$

Solution Taking the Aboodh transform of equation of both sides of (4.23)

$$u\mathcal{A}\{\phi(y, t)\} - \frac{\phi(y, 0)}{u} = \mathcal{A}\left\{\frac{\partial^2 \phi(y, t)}{\partial y^2} + 6(1 - \phi)\phi\right\} \quad (4.25)$$

$$u\mathcal{A}\{\phi(y, t)\} = \frac{\phi(y, 0)}{u} + \mathcal{A}\left\{\frac{\partial^2 \phi(y, t)}{\partial y^2} + 6(1 - \phi)\phi\right\} \quad (4.26)$$

Multiplying equation (4.26) by $\frac{1}{u}$. Then we have,

$$\mathcal{A}\{\phi(y, t)\} = \frac{\phi(y, 0)}{u^2} + \frac{1}{u}\mathcal{A}\left\{\frac{\partial^2 \phi(y, t)}{\partial y^2} + 6(1 - \phi)\phi\right\} \quad (4.27)$$

Taking the Aboodh inverse transform of (4.27)

$$\mathcal{A}\{\phi(y, t)\} = \mathcal{A}^{-1}\left[\frac{e^y}{u^2}\right] + \mathcal{A}^{-1}\left[\frac{1}{u}\mathcal{A}\left\{\frac{\partial^2 \phi(y, t)}{\partial y^2} + 6(1 - \phi)\phi\right\}\right] \quad (4.28)$$

Applying HPM technique to equation (4.28). Thus, we get

$$\sum_{m=0}^{\infty} \phi_{m+1}(y, t) = e^y + p \left(\mathcal{A}^{-1} \left\{ \frac{1}{u} \mathcal{A}^{-1} \left[\frac{\partial^2}{\partial y^2} \sum_{m=0}^{\infty} p^m \phi_m(y, t) + 6 \sum_{m=0}^{\infty} p^m \phi_m(y, t) - 6 \sum_{m=0}^{\infty} p^m \phi_m^2 \right] \right\} \right) \quad (4.29)$$

Applying He's polynomial to simplify the nonlinear terms,

$$\sum_{m=0}^{\infty} p^m \phi_m^2(\phi) = \sum_{m=0}^{\infty} p^m H_m(\phi) \quad (4.30)$$

$$\sum_{m=0}^{\infty} \phi_m(y, t) = \phi_0(y, t) + p\phi_1(y, t) + p^2\phi_2(y, t) + \dots \quad (4.31)$$

Substituting (4.30) and (4.31) into equation (4.29), we have

$$\sum_{m=0}^{\infty} \phi_{m+1}(y, t) = e^y + p \left(\mathcal{A}^{-1} \left\{ u^2 \mathcal{A} \left[\frac{\partial^2}{\partial y^2} \sum_{m=0}^{\infty} p^m \phi_m(y, t) + 6 \sum_{m=0}^{\infty} p^m \phi_m(y, t) - 6 \sum_{m=0}^{\infty} p^m H_m(\phi) \right] \right\} \right) \quad (4.32)$$

Using HPM, the series becomes

$$\begin{aligned} \sum_{m=0}^{\infty} \phi_{m+1}(y, t) = e^y + p \left(\mathcal{A}^{-1} \left\{ u^2 \mathcal{A} \left[\left(\frac{\partial^2}{\partial y^2} \sum_{m=0}^{\infty} p^m \phi_m(y, t) + 6 \sum_{r=0}^{\infty} p^r \phi_r(y, t) - 6 \sum_{r=0}^{\infty} H_r(\phi) \right) \right. \right. \right. \\ \left. \left. \left. + p \left(\frac{\partial^2}{\partial y^2} \sum_{m=0}^{\infty} p^m \phi_m(y, t) + 6 \sum_{m=0}^{\infty} p^m \phi_m(y, t) - 6 \sum_{m=0}^{\infty} H_m(\phi) \right) + \dots \right] \right\} \right) \end{aligned} \quad (4.33)$$

Comparing the respective powers of p of equation (4.33), the following approximations are obtained.

$$\phi_0(y, t) = \phi(y, 0) = e^y \quad (4.34)$$

$$H_0(\phi) = \phi_0 \phi_1 \quad (4.35)$$

$$\begin{aligned} \phi_1(y, t) &= \mathcal{A}^{-1} \left\{ \frac{1}{u} \mathcal{A} \left(\frac{\partial^2 \phi_0}{\partial y^2} + 6\phi_0 - 6H_0(\phi) \right) \right\} \\ &= \mathcal{A}^{-1} \left\{ \frac{1}{u} \mathcal{A} (e^y + 6e^y - e^{2y}) \right\} \\ &= \mathcal{A}^{-1} \left\{ \frac{1}{u} (e^y + 6e^y - 6ve^{2y}) \right\} \\ &= \mathcal{A}^{-1} \left\{ \frac{1}{u} (7e^y - 6e^{2y}) \right\} \\ &= 7te^y - 6te^{2y} \\ &= (7 - 6e^y) te^y \end{aligned} \quad (4.36)$$

The approximate solution to equation (4.23) is given as

$$\sum_{r=0}^{\infty} \phi(y, t) = e^y + (7 + 6e^y) te^y + \dots \quad (4.37)$$

Equation (4.38) is the same as the solution obtained in [25]

Example 4.3. Considering the one dimensional Fisher's equation of the form [25]

$$\frac{\partial \phi}{\partial t}(y, t) - \frac{\partial^2 \phi}{\partial y^2} - 6\phi(y, t) [1 - \phi(y, t)] = 0 \quad (4.38)$$

With the initial condition

$$\phi(y, 0) = \frac{1}{(1 + e^x)^2} \quad (4.39)$$

Solution

Equation (4.39) can also be written as

$$\frac{\partial \phi}{\partial t}(y, t) = \frac{\partial^2 \phi}{\partial y^2} + 6\phi(y, t)[1 - \phi(y, t)] \quad (4.40)$$

Taking the Aboodh transform of equation (4.41), we get

$$v\mathcal{A}\{\phi(y, t)\} - \frac{\phi(y, 0)}{u} = \mathcal{A}\left\{ \frac{\partial^2 \phi}{\partial y^2} + 6\phi(y, t)[1 - \phi(y, t)] \right\} \quad (4.41)$$

Multiplying equation (4.42) by $\frac{1}{u}$. Then, the equation becomes

$$u\mathcal{A}\{\phi(y, t)\} = \frac{\phi(y, 0)}{u^2} + \frac{1}{u}\mathcal{A}\left\{\frac{\partial^2\phi}{\partial y^2} + 6\phi(y, t)[1 - \phi(y, t)]\right\} \quad (4.43)$$

Taking the Aboodh inverse transform of both sides of equation (4.43). Thus,

$$\phi(y, t) = \mathcal{A}\left\{\frac{\phi(y, 0)}{u^2}\right\} + \mathcal{A}^{-1}\left\{\frac{1}{u}\mathcal{A}\left[\frac{\partial^2\phi}{\partial y^2} + 6\phi(y, t)[1 - \phi(y, t)]\right]\right\} \quad (4.44)$$

Applying the HPM technique to assume the series of solution of the function $\phi(y, t)$. Then, we obtain

$$\phi(y, t) = \sum_{m=0}^{\infty} \phi_m(y, t) \quad (4.45)$$

Using equation (4.45) in equation (4.44), we get

$$\sum_{m=0}^{\infty} \phi_{m+1}(y, t) = \frac{1}{(1+e^y)^2} + p \left(\mathcal{A}^{-1} \left\{ \frac{1}{u} \mathcal{A} \left[\frac{\partial^2}{\partial y^2} \sum_{m=0}^{\infty} p^m \phi_m(y, t) + 6 \sum_{m=0}^{\infty} p^m \phi_m(y, t) - 6 \sum_{m=0}^{\infty} p^m \phi_m^2 \right] \right\} \right) \quad (4.46)$$

Applying He's polynomial to simplify the nonlinear terms in equation (4.46),

$$\sum_{m=0}^{\infty} p^m \phi_m^2 = \sum_{m=0}^{\infty} p^m H_m^2(\phi) \quad (4.47)$$

$$\sum_{m=0}^{\infty} \phi_m(y, t) = \phi_0(y, t) + p\phi_1(y, t) + p^2\phi_2(y, t) + \dots \quad (4.48)$$

Substituting (4.47) and (4.48) into equation (4.46), we have

$$\sum_{m=0}^{\infty} \phi_{m+1}(y, t) = \frac{1}{(1+e^y)^2} + p \left(\mathcal{A}^{-1} \left\{ \frac{1}{u} \mathcal{A} \left[\frac{\partial^2}{\partial y^2} \sum_{m=0}^{\infty} p^m \phi_m(y, t) + 6 \sum_{m=0}^{\infty} p^m \phi_m(y, t) - 6 \sum_{m=0}^{\infty} p^m H_m(\phi) \right] \right\} \right) \quad (4.49)$$

$$\begin{aligned} \sum_{m=0}^{\infty} \phi_{m+1}(y, t) = & \frac{1}{(1+e^y)^2} + p \left(\mathcal{A}^{-1} \left\{ \frac{1}{u} \mathcal{A} \left[\left(\frac{\partial^2}{\partial y^2} \sum_{m=0}^{\infty} \phi_0(y, t) + 6 \sum_{m=0}^{\infty} \phi_0(y, t) - 6 \sum_{m=0}^{\infty} H_m(\phi) \right) \right. \right. \right. \\ & \left. \left. \left. + p \left(\frac{\partial^2}{\partial y^2} \sum_{m=0}^{\infty} \phi_1(y, t) + 6 \sum_{m=0}^{\infty} \phi_1(y, t) - 6 \sum_{m=0}^{\infty} H_m(\phi) \right) + \dots \right] \right\} \right) \quad (4.50) \end{aligned}$$

Using HPM, the series becomes
Comparing the respective powers of p of equation (4.50), the following approximations are obtained.

$$\phi_0(y, t) = \phi \quad (4.51)$$

$$(y, 0) = \frac{1}{(1 + e^y)^2}$$

$$H_0(\phi) = \phi_0 \phi_1$$

$$\begin{aligned} \phi_1(y, t) &= \mathcal{A}^{-1} \left\{ \frac{1}{u} \mathcal{A} \left(\frac{\partial^2 \phi_0}{\partial y^2} + 6\phi_0 - 6H_0(\phi) \right) \right\} \\ &= \frac{10e^y t}{(1 + e^y)^3} \end{aligned} \quad (4.52)$$

$$\begin{aligned} \phi_2(y, t) &= \mathcal{A}^{-1} \left\{ \frac{1}{u} \mathcal{A} \left(\frac{\partial^2 \phi}{\partial y^2} + 6\phi_1 - 6H_1(\phi) \right) \right\} \\ &= \frac{25e^y (-1 + 2e^y) t^2}{(1 + e^y)^4} \end{aligned} \quad (4.53)$$

The approximate solution to equation (4.39) is given as

$$\sum_{m=0}^{\infty} \phi_{m+1}(y, t) = \frac{1}{(1 + e^y)^2} + \frac{10e^y t}{(1 + e^y)^3} + \frac{25e^y (-1 + 2e^y) t^2}{(1 + e^y)^4} + \dots \quad (4.54)$$

Equation (4.54) is the same as the exact solution obtained in [25]

5. CONCLUSION

In this work, one dimensional Fisher's problem was solved by using the combination of Aboodh transform and homotopy perturbation method (HPM). The outcomes demonstrate the effectiveness and potency of the AHPM approach in locating both precise and approximate solutions to nonlinear differential equations. When compared to previous methods, AHPM provides very accurate numerical solutions for nonlinear problems because it yields an efficient recurring relation for solving Fishers Type equations. According to this approach, the answer is found by adding an infinite series that quickly converges to the precise answers.

REFERENCES

1. K. Aboodh, *Homotopy perturbation method and Aboodh transform for solving nonlinear partial differential equations*. Theory of Approximation and Applications, **11**(1), (2017) pp.1-12.
2. D. Ağırseven, and T. Öziş, *An analytical study for Fisher type equations by using homotopy perturbation method*. Computers & Mathematics with Applications, **60**(3), (2010), pp.602-609. <http://dx.doi.org/10.1016/j.camwa.2010.05.006>
3. H. O. Al-Humedi, & A. H. Ali *Application of Adomian decomposition method to solve Fisher's equation*. Journal of Basrah Researches (Sciences), **35**, (2008), pp.23-32.
4. A. A. Aswhad, & A. F. Jaddoa, *The approximate solution of newell whitehead segel and fisher equations using the adomian decomposition method* (2016) <http://dx.doi.org/10.5281/zenodo.50342>

5. N. Bildik & A., Konuralp, *The use of variational iteration method, differential transform method and Adomian decomposition method for solving different types of nonlinear partial differential equations*. International Journal of Nonlinear Sciences and Numerical Simulation, **7**(1), (2006) 65-70. <http://dx.doi.org/10.1515/IJNSNS.2006.7.1.65>
6. V. Chandraker, A., Awasthi, & S. Jayaraj, *A numerical treatment of Fisher equation*. Procedia Engineering, **127**, (2015), pp.1256-1262. <https://doi.org/10.1016/j.proeng.2015.11.481>
7. Z. Z. Ganji, D. D., Ganji, H., Jafari, & M., Rostamian, *Application of the homotopy perturbation method to coupled system of partial differential equations with time fractional derivatives*. Topological methods in nonlinear analysis. **31** (2008) 341-348.
8. B. H. Gilding, & R. Kesner, *A Fisher/KPP-type equation with density-dependent diffusion and convection: travelling-wave solutions*. Journal of Physics A: Mathematical and General, **38**(15), (2005), p.3367. <https://doi.org/10.1088/0305-4470/38/15/009>
9. A. Grafen, *A theory of Fisher's reproductive value*. Journal of mathematical biology, **53**, (2006) pp.15-60.
10. V. G. Gupta, & S. Gupta, *Applications of homotopy analysis transform method for solving various nonlinear equations*. World Applied Sciences Journal, **18**(12), (2012) 1839-1846. <http://dx.doi.org/10.5829/idosi.wasj.2012.18.12.2700>
11. J. H. He, *A coupling method of a homotopy technique and a perturbation technique for nonlinear problems*. International journal of non-linear mechanics, **35**(1), (2000), pp.37-43. [http://dx.doi.org/10.1016/S0020-7462\(98\)00085-7](http://dx.doi.org/10.1016/S0020-7462(98)00085-7)
12. S. A. Levin, B. Grenfell, A. Hastings, A. S. and Perelson, *Mathematical and computational challenges in population biology and ecosystems science*. Science, **275**(5298), (1997), pp.334-343. <https://doi.org/10.1126/science.275.5298.334>
13. M. I. Liaqat, A. Khan, M. A. Alam, and M. K. Pandit, *A highly accurate technique to obtain exact solutions to time-fractional quantum mechanics problems with zero and nonzero trapping potential*. Journal of Mathematics, **2022**, (2022) pp.1-20. <http://dx.doi.org/10.3390/sym15030744>
14. M. I., Liaqat, A., Khan, M. A., Alam, M. K., Pandit, S. Etemad, & Rezapour, *Approximate and closed-form solutions of Newell-Whitehead-Segel equations via modified conformable Shehu transform decomposition method*. Mathematical Problems in Engineering (2022) , <http://dx.doi.org/10.1155/2022/6752455>
15. M. Madani, M. Fathizadeh, Y., Khan, & Yildirim, A. *On the coupling of the homotopy perturbation method and Laplace transformation*. Mathematical and Computer Modelling, **53**(9-10), (2011) 1937-1945. <http://dx.doi.org/10.1016/j.mcm.2011.01.023>
16. K. Manimegalai, C. F. Zephania Segar, P. K. Bera, P. Bera, S. K. Das, and T. Sil, 2019. *Study of strongly nonlinear oscillators using the Aboodh transform and the homotopy perturbation method*. The European Physical Journal Plus, **134**, (2019), pp.1-10. <http://dx.doi.org/10.1140/epjp/i2019-12824-6>
17. S. S. Nourazar, M. Soori, & A. Nazari-Golshan, *On the exact solution of Newell-Whitehead-Segel equation using the homotopy perturbation method* (2015).
18. O. Olabanjo, & A. Wusu, *Dynamics of sustainable fisheries: a mathematical approach using Lotka-Volterra equations*. Annals of Mathematics and Computer Science, **19**, (2023) 59-67, <https://doi.org/10.56947/amcs.v19.225>
19. D. Olmos, & B. D. Shizgal, *A pseudospectral method of solution of Fisher's equation*. Journal of Computational and Applied Mathematics, **193**(1), (2006), pp.219-242. <https://doi.org/10.1016/j.cam.2005.06.028>
20. Olubanwo, O. O., Odetunde, O. S., & Talabi, A. T. *Aboodh homotopy perturbation method of solving burgers equation*. Asian Journal of Applied Sciences, **7**(2), (2019), <https://doi.org/10.24203/ajas.v7i2.5794>
21. O. O. Olubanwo, & O. S. Odetunde, & T. Adetoro, (2019). *Aboodh Homotopy Perturbation Method of solving Burgers Equation*. Asian Journal of Applied Sciences. **7**, (2019), <https://doi.org/10.24203/ajas.v7i2.5794>.

22. O. O. Olubanwo, J. T. Adepoju, A. S. Ajani, & E. A. Shobowale. *Application of Mohand transform coupled with homotopy perturbation method to solve Newel-White-Segel equation*. Annals of Mathematics and Computer Science, **21** (2024), 162-180, <https://doi.org/10.56947/amcs.v21.267>
23. Olubanwo, O.O., and Odetunde, O. S. "Laplace Homotopy Perturbation Perturbation Method of Solving Nonlinear Partial Differential Equations." Annals. Computer Science Series **17**, no. **2** (2019).
24. Olubanwo, O.O., Talabi, A.T., Odetunde, O.S. and Ogunwobi, Z.O., Modified Aboodh Homotopy Perturbation Method For Solving Nonlinear Gas Dynamic Equation. LAUTECH Journal of Engineering and Technology, **13**(1), (2019), pp.1-12
25. K. Pakhtunkhwa, & Correspondence, Pakistan & S. Kamal & A. Arshad & H. Laila & H. Ali., *Analytical solution of General Fisher's Equation by using Laplace Adomian decomposition method.*, Journal of Pure and Applied Mathematics, **2021**, (2021).
26. N. Parekh, and S., Puri, S., *A new numerical scheme for the Fisher equation*. Journal of physics A: Mathematical and general, **23**(21), (1990), p.L1085. <http://dx.doi.org/10.1088/0305-4470/23/21/003>
27. P., Pue-On, *Laplace Adomian decomposition method for solving Newell-Whitehead-Segel equation*. Applied Mathematical Sciences, **7**(132), (2013), 6593-6600. <http://dx.doi.org/10.12988/ams.2013.310603>
28. H. Tao, N. Anjum, and Y. J. Yang, *The Aboodh transformation-based homotopy perturbation method: new hope for fractional calculus*. Frontiers in Physics, **11**, (2023) p.1168795. <http://dx.doi.org/10.3389/fphy.2023.1168795>.

¹ DEPARTMENT OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCE, OLABISI ONABANJO UNIVERSITY, AGO-IWOYE, OGUN STATE, NIGERIA

Email address: olubanwo.oludapo@oouagoiwoye.edu.ng; adepojujulius58@gmail.com; ajani.abiodun@oouagoiwoye.edu.ng; idowusunday288@gmail.com